



CHICK STARTER GROWER KRUMBLE (MEDICATED)

Complete feed for starting and growing replacement chickens.

For the development of active immunity to coccidiosis and for increased rate of weight gain and improved feed efficiency in replacement chickens.

ACTIVE DRUG INGREDIENTS:

Amprolium.....0.0125%
Bacitracin Methylene Disalicylate..... 50 g/ton

GUARANTEED ANALYSIS:

Crude Protein, min.....18.00%
Lysine, min.....0.95%
Methionine, min.....0.40%
Crude Fat, min.....3.00%
Crude Fiber, max.....6.50%
Calcium (Ca), min.....1.00%, max.....1.50%
Phosphorus (P), min.....0.68%
Salt (NaCl), min.....0.05%, max.....0.55%
Vitamin A, min.....4,800 IU/Lb.
Vitamin D3, min..... 1770 IU/lb.
Vitamin E, min..... 35 IU/lb.

INGREDIENTS:

Corn, Soybean Meal, Wheat Middling, Rice Bran, Alfalfa Meal Dehydrated, Animal Protein Products, Calcium Carbonate, Monocalcium/Dicalcium Phosphate, Salt, DL-Methionine, L-Lysine, Saccharomyces cerevisiae, Choline Chloride, Ferrous Sulfate, Manganese Hydroxychloride, Manganese Amino Acid Complex, Zinc Hydroxychloride, Zinc Amino Acid Complex, Vitamin E Supplement, Niacin Supplement, Sodium Selenite, Vitamin A Supplement, Basic Copper Chloride, Copper Amino Acid Complex, Dried Bacillus Licheniformis Fermentation Product, Dried Bacillus Subtilis Fermentation Product, Dried Trichoderma Reesei Fermentation Products (Phytase), Sodium Phytate, Riboflavin Supplement, D-Calcium Pantothenate, Folic Acid, Vitamin D3 Supplement, Thiamine Mononitrate, Menadione Dimethylpyrimidinol Bisulfite, Ethylenediamine Dihydroiodide, Pyridoxine Hydrochloride, Biotin, Vitamin B12 Supplement, Natural & Artificial Flavor Ingredients.

FEEDING DIRECTIONS:

Feed continuously as the sole ration under conditions of slight exposure to coccidiosis up to 5 weeks of age, from 5 to 8 weeks of age or over 8 weeks of age.

For raising layers after 18 weeks of age, change to either Big V 20% Egg Pellets and Scratch grain or Big V 16% All mash Laying pellet. Keep plenty of fresh clean water available at all times. As an option, feed oyster shell and grit free choice. Results may vary depending upon feed intake, environmental conditions and the quality of management practices.

WARNING: Use as sole source of Amprolium.

CAUTION: For replacement chicken only. Do not use AMPROLIUM in feeds containing bentonite.

Ruminant Meat and Bone Meal Free!

Contains Natural Botanical Products!



**MANUFACTURED BY
BIG V FEEDS, INC.
MCALISTER, OKLAHOMA 74502**
www.bigvfeeds.com

Net Wt. 50 Lbs. 22.68 kg.



CHICK STARTER GROWER KRUMBLE (MEDICATED)

Complete feed for starting and growing replacement chickens.

For the development of active immunity to coccidiosis and for increased rate of weight gain and improved feed efficiency in replacement chickens.

ACTIVE DRUG INGREDIENTS:

Amprolium.....0.0125%
Bacitracin Methylene Disalicylate..... 50 g/ton

GUARANTEED ANALYSIS:

Crude Protein, min.....18.00%
Lysine, min.....0.95%
Methionine, min.....0.40%
Crude Fat, min.....3.00%
Crude Fiber, max.....6.50%
Calcium (Ca), min.....1.00%, max.....1.50%
Phosphorus (P), min.....0.68%
Salt (NaCl), min.....0.05%, max.....0.55%
Vitamin A, min.....4,800 IU/Lb.
Vitamin D3, min..... 1770 IU/lb.
Vitamin E, min..... 35 IU/lb.

INGREDIENTS:

Corn, Soybean Meal, Wheat Middling, Rice Bran, Alfalfa Meal Dehydrated, Animal Protein Products, Calcium Carbonate, Monocalcium/Dicalcium Phosphate, Salt, DL-Methionine, L-Lysine, Saccharomyces cerevisiae, Choline Chloride, Ferrous Sulfate, Manganese Hydroxychloride, Manganese Amino Acid Complex, Zinc Hydroxychloride, Zinc Amino Acid Complex, Vitamin E Supplement, Niacin Supplement, Sodium Selenite, Vitamin A Supplement, Basic Copper Chloride, Copper Amino Acid Complex, Dried Bacillus Licheniformis Fermentation Product, Dried Bacillus Subtilis Fermentation Product, Dried Trichoderma Reesei Fermentation Products (Phytase), Sodium Phytate, Riboflavin Supplement, D-Calcium Pantothenate, Folic Acid, Vitamin D3 Supplement, Thiamine Mononitrate, Menadione Dimethylpyrimidinol Bisulfite, Ethylenediamine Dihydroiodide, Pyridoxine Hydrochloride, Biotin, Vitamin B12 Supplement, Natural & Artificial Flavor Ingredients.

FEEDING DIRECTIONS:

Feed continuously as the sole ration under conditions of slight exposure to coccidiosis up to 5 weeks of age, from 5 to 8 weeks of age or over 8 weeks of age.

For raising layers after 18 weeks of age, change to either Big V 20% Egg Pellets and Scratch grain or Big V 16% All mash Laying pellet. Keep plenty of fresh clean water available at all times. As an option, feed oyster shell and grit free choice. Results may vary depending upon feed intake, environmental conditions and the quality of management practices.

WARNING: Use as sole source of Amprolium.

CAUTION: For replacement chicken only. Do not use AMPROLIUM in feeds containing bentonite.

Ruminant Meat and Bone Meal Free!

Contains Natural Botanical Products!



**MANUFACTURED BY
BIG V FEEDS, INC.
MCALISTER, OKLAHOMA 74502**
www.bigvfeeds.com

Net Wt. 50 Lbs. 22.68 kg.

