



15% RANGE MEAL; CTC/IGR

(MEDICATED)

For beef cattle on pasture.



CAUTION:

Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella spp.* susceptible to Chlortetracycline. Control of active infection of anaplasmosis caused by *Anaplasma Marginale* susceptible to chlortetracycline.

ACTIVE DRUG INGREDIENT:

Chlortetracycline..... 750 gms/ton

Insect growth regulator for continuous feeding during the fly season to prevent the breeding of horn flies in the manure of treated cattle.

ACTIVE INGREDIENT:

(S)-Methoprene (CAS#65733-16-6)..... 0.0017%
Other Ingredients.....99.9983%
Total 100.000%

GUARANTEED ANALYSIS:

Crude Protein, min.....15.00%
Crude Fat, min..... 3.50%
Crude Fiber, max.....8.00%
Calcium (Ca), min..... 2.25%, max..... 2.75%
Phosphorus (P), min..... 0.40%
Salt (NaCl), min.....25.00%, max.....29.00%
Potassium (K), min.....0.75%
Vitamin A, min.....20,000 IU/lb.

INGREDIENTS:

Processed Grain By-Products, Salt, Plant Protein Products, Grain Products, Roughage Products, Calcium Carbonate, Molasses Products, Zinc Hydroxychloride, Potassium Chloride, Magnesium Oxide, Basic Copper Chloride, Vitamin A Supplement, Manganese Hydroxychloride, Sodium Selenite, Vitamin D3 Supplement, Vitamin E Supplement, Cobalt Carbonate, Ferrous Sulfate, Ethylenediamine Dihydroiodide.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Each pound of this feed contains 375 mg. of chlortetracycline. For control of **bacterial pneumonia** associated with shipping fever complex; feed beef cattle at a rate to provide 350 mg/head/day (0.93 lb. of feed). For control of **active infection of anaplasmosis**; beef cattle under 700 lbs. feed 350 mg./head/day (0.93 lbs. of feed). For beef cattle over 700 lbs. feed 0.5 mg./lb. of body weight daily. (Example 1000 lb. animal needs 500 mgs (1.34 lbs. of feed)/day; a 1500 lb. animal needs 750 mgs (2 lbs. of feed)/day.)

These feeding rates will provide 23 to 45.4 mg of (S)-methoprene per 100 lbs. body weight per month. Introduction of this product after adult horn fly infestation is established will require treatment of cattle with adulticides if elimination of the adult fly population is desired. Provide clean fresh water at all times and a good roughage source.

ENVIRONMENTAL HAZARDS: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

CAUTION: Contains added copper, consumption of this product by sheep may result in copper toxicity.

WARNING: A withdraw period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. **Storage:** Store this product in a cool, dry place. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Handling:** Completely empty bag, dispose bag in a sanitary landfill or by incineration or if allowed by state and local authorities by burning. If burned, stay out of smoke.

MANUFACTURED BY

BIG Y FEEDS, INC.

MCALISTER, OK 74502

WWW.BIGYFEEDS.COM

Net Wt. 50 LBS. (22.68 KGS.)

0284



15% RANGE MEAL; CTC/IGR

(MEDICATED)

For beef cattle on pasture.



CAUTION:

Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella spp.* susceptible to Chlortetracycline. Control of active infection of anaplasmosis caused by *Anaplasma Marginale* susceptible to chlortetracycline.

ACTIVE DRUG INGREDIENT:

Chlortetracycline..... 750 gms/ton

Insect growth regulator for continuous feeding during the fly season to prevent the breeding of horn flies in the manure of treated cattle.

ACTIVE INGREDIENT:

(S)-Methoprene (CAS#65733-16-6)..... 0.0017%
Other Ingredients.....99.9983%
Total 100.000%

GUARANTEED ANALYSIS:

Crude Protein, min.....15.00%
Crude Fat, min..... 3.50%
Crude Fiber, max.....8.00%
Calcium (Ca), min..... 2.25%, max..... 2.75%
Phosphorus (P), min..... 0.40%
Salt (NaCl), min.....25.00%, max.....29.00%
Potassium (K), min.....0.75%
Vitamin A, min.....20,000 IU/lb.

INGREDIENTS:

Processed Grain By-Products, Salt, Plant Protein Products, Grain Products, Roughage Products, Calcium Carbonate, Molasses Products, Zinc Hydroxychloride, Potassium Chloride, Magnesium Oxide, Basic Copper Chloride, Vitamin A Supplement, Manganese Hydroxychloride, Sodium Selenite, Vitamin D3 Supplement, Vitamin E Supplement, Cobalt Carbonate, Ferrous Sulfate, Ethylenediamine Dihydroiodide.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Each pound of this feed contains 375 mg. of chlortetracycline. For control of **bacterial pneumonia** associated with shipping fever complex; feed beef cattle at a rate to provide 350 mg/head/day (0.93 lb. of feed). For control of **active infection of anaplasmosis**; beef cattle under 700 lbs. feed 350 mg./head/day (0.93 lbs. of feed). For beef cattle over 700 lbs. feed 0.5 mg./lb. of body weight daily. (Example 1000 lb. animal needs 500 mgs (1.34 lbs. of feed)/day; a 1500 lb. animal needs 750 mgs (2 lbs. of feed)/day.)

These feeding rates will provide 23 to 45.4 mg of (S)-methoprene per 100 lbs. body weight per month. Introduction of this product after adult horn fly infestation is established will require treatment of cattle with adulticides if elimination of the adult fly population is desired. Provide clean fresh water at all times and a good roughage source.

ENVIRONMENTAL HAZARDS: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

CAUTION: Contains added copper, consumption of this product by sheep may result in copper toxicity.

WARNING: A withdraw period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. **Storage:** Store this product in a cool, dry place. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Handling:** Completely empty bag, dispose bag in a sanitary landfill or by incineration or if allowed by state and local authorities by burning. If burned, stay out of smoke.

MANUFACTURED BY

BIG Y FEEDS, INC.

MCALISTER, OK 74502

WWW.BIGYFEEDS.COM

Net Wt. 50 LBS. (22.68 KGS.)

0284