



WILDLIFE AND ZOO RUMINANTS FEED with Safe-Guard® (Apple Flavor) FENBENDAZOLE TYPE C MEDICATED FEED INDICATIONS

For the treatment and control of: **Stomach worms:** small stomach worms (*Trichostrongylus* spp.), barberpole worm (*Haemonchus* spp.); **Intestinal worms:** thread necked intestinal worm (*Nematodirus* spp.), and whipworm (*Trichuris* spp.) for the following animal species:

Ruminants— Subfamily antilopinae:

Persain gazelles (*Gazella subgutturosa*)
Addra gazelle (*Gazella dama ruficollis*)
Slenderhorn gazelle (*Gazella leptoceros*)
Kenya impala (*Aepyceros melampus redilis*)
Roosevelt's gazelle (*Gazella granti roosevelti*)
Indian Blackbuck (*Antelope cervicapra*)
Mhor gazelle (*Gazella dama mhor*)
Thomson's gazelles (*Gazella thomsoni*)

Ruminants— Subfamily Hippotraginae:

Addax (*Addax nasomaculatus*)
Angolan roan antelope (*Hippotragus equinus cottoni*)
Fringed-ear oryx (*Oryx gazelle callotis*)
Arabian oryx (*Oryx leucoryx*)
Ruminants— Subfamily caprinae:
Armenian mouflon (*Ovis orientalis gmelini*)
Russian saiga (*Saiga tatarica*)

ACTIVE DRUG INGREDIENT:

Fenbendazole.....(114 mg/lb.).... 228 g/ton

GUARANTEED ANALYSIS:

Crude Protein, min. 16.00%
Crude Fat, min. 4.00%
Crude Fiber, max..... 12.00%
Calcium (Ca), min. 1.00%, max.....1.50%
Phosphorus (P), min.0.70%
Salt (NaCl), min.0.75%, max.....1.25%
Potassium (K), min.1.00%

INGREDIENTS:

Processed Grain By-Products, Grain Products, Plant Protein Products, Roughage Products, Magnesium Mica, Calcium Carbonate, Salt, Natural and Artificial Flavors.

FEEDING DIRECTIONS:

Feed at a rate to provide 1.14 mg of Fenbendazole per pound of body weight per day for (3) consecutive days. Type C feeds containing Fenbendazole can be fed to adult and young animals. No prior withdrawal of feed or water is necessary.

The following example chart gives the feeding rate of this Type C Fenbendazole medicated feed based on total group animal body weight to provide 1.14 mg/lb./day dosage. **Feed daily for (3) days.**

Retreatment may be required in six weeks, depending upon conditions of continued exposure to parasites, condition of treated animals and ambient temperatures.

weight	100 lbs.	500 lbs.	1,000 lbs.	3,000 lbs.	5,000 lbs.
Feed	1.0 lb.	5.0 lbs.	10.0 lbs.	30.0 lbs.	50.0 lbs.

► RESIDUE WARNING ◀

Do not use 14 days before or during hunting season.

OTHER WARNINGS

Parasite resistance may develop to any dewormer. All dewormers require accurate dosing for best results. Following the use of any dewormer, effectiveness of treatment should be monitored. A decrease of effectiveness over time may indicate the development of resistance to the dewormer administered. The parasite management plan should be adjusted accordingly based on regular monitoring.

Consult veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Approved by FDA, under NADA# 131-675

Safe-Guard is a registered trademark of Intervet International B.V.



WILDLIFE AND ZOO RUMINANTS FEED with Safe-Guard® (Apple Flavor) FENBENDAZOLE TYPE C MEDICATED FEED INDICATIONS

For the treatment and control of: **Stomach worms:** small stomach worms (*Trichostrongylus* spp.), barberpole worm (*Haemonchus* spp.); **Intestinal worms:** thread necked intestinal worm (*Nematodirus* spp.), and whipworm (*Trichuris* spp.) for the following animal species:

Ruminants— Subfamily antilopinae:

Persain gazelles (*Gazella subgutturosa*)
Addra gazelle (*Gazella dama ruficollis*)
Slenderhorn gazelle (*Gazella leptoceros*)
Kenya impala (*Aepyceros melampus redilis*)
Roosevelt's gazelle (*Gazella granti roosevelti*)
Indian Blackbuck (*Antelope cervicapra*)
Mhor gazelle (*Gazella dama mhor*)
Thomson's gazelles (*Gazella thomsoni*)

Ruminants— Subfamily Hippotraginae:

Addax (*Addax nasomaculatus*)
Angolan roan antelope (*Hippotragus equinus cottoni*)
Fringed-ear oryx (*Oryx gazelle callotis*)
Arabian oryx (*Oryx leucoryx*)
Ruminants— Subfamily caprinae:
Armenian mouflon (*Ovis orientalis gmelini*)
Russian saiga (*Saiga tatarica*)

ACTIVE DRUG INGREDIENT:

Fenbendazole.....(114 mg/lb.).... 228 g/ton

GUARANTEED ANALYSIS:

Crude Protein, min. 16.00%
Crude Fat, min. 4.00%
Crude Fiber, max..... 12.00%
Calcium (Ca), min. 1.00%, max.....1.50%
Phosphorus (P), min.0.70%
Salt (NaCl), min.0.75%, max.....1.25%
Potassium (K), min.1.00%

INGREDIENTS:

Processed Grain By-Products, Grain Products, Plant Protein Products, Roughage Products, Magnesium Mica, Calcium Carbonate, Salt, Natural and Artificial Flavors.

FEEDING DIRECTIONS:

Feed at a rate to provide 1.14 mg of Fenbendazole per pound of body weight per day for (3) consecutive days. Type C feeds containing Fenbendazole can be fed to adult and young animals. No prior withdrawal of feed or water is necessary.

The following example chart gives the feeding rate of this Type C Fenbendazole medicated feed based on total group animal body weight to provide 1.14 mg/lb./day dosage. **Feed daily for (3) days.**

Retreatment may be required in six weeks, depending upon conditions of continued exposure to parasites, condition of treated animals and ambient temperatures.

weight	100 lbs.	500 lbs.	1,000 lbs.	3,000 lbs.	5,000 lbs.
Feed	1.0 lb.	5.0 lbs.	10.0 lbs.	30.0 lbs.	50.0 lbs.

► RESIDUE WARNING ◀

Do not use 14 days before or during hunting season.

OTHER WARNINGS

Parasite resistance may develop to any dewormer. All dewormers require accurate dosing for best results. Following the use of any dewormer, effectiveness of treatment should be monitored. A decrease of effectiveness over time may indicate the development of resistance to the dewormer administered. The parasite management plan should be adjusted accordingly based on regular monitoring.

Consult veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Approved by FDA, under NADA# 131-675

Safe-Guard is a registered trademark of Intervet International B.V.



MANUFACTURED BY
BIG Y FEEDS, INC., MCALISTER, OK 74502
Net Wt. 25 LBS. (11.34 KG.)



MANUFACTURED BY
BIG Y FEEDS, INC., MCALISTER, OK 74502
Net Wt. 25 LBS. (11.34 KG.)