



BIG MAC BREEDER CUBE

WITH SAFEGUARD® DEWORMER

Type C Medicated Feed

For beef cattle on pasture.

INDICATIONS

For the treatment and control of: **Lungworms:** Adult *Dictyocaulus viviparus*; **Stomach worms:** Adult brown stomach worms (*Ostertagia ostertagi*), Adult and fourth stage larvae barberpole worms (*Haemonchus contortus*), fourth stage larvae barberpole worms (*H. placei*), and Adult and fourth stage larvae small stomach worms (*Trichostrongylus axei*); **Intestinal worms** (Adult and fourth stage larvae): Hookworms (*Bunostomum phlebotomum*), thread-necked intestinal worms (*Nematodirus helvetianus*), small intestinal worms (*Cooperia punctata* & *C. oncophora*), bankrupt worms (*Trichostrongylus colubriformis*), and nodular worms (*Oesophagostomum radiatum*).

ACTIVE DRUG INGREDIENT:

Fenbendazole.....454 mg/lb.

GUARANTEED ANALYSIS:

Crude Protein, min.20.00%
Crude Fat, min.2.60%
Crude Fiber, max.10.00%
Calcium (Ca), min.0.75%, max.1.25%
Phosphorus (P), min.0.75%
Salt (NaCl), min.0.75%, max.1.25%
Potassium (K), min.1.00%
Vitamin A, min.20,000 IU/Lb.



INGREDIENTS:

Processed Grain By-Products, Plant Protein Products, Roughage Products, Grain Products, Molasses Products, Magnesium Mica, Calcium Carbonate, Salt, Vitamin A Supplement, Vitamin D Supplement.

FEEDING DIRECTIONS:

Feed as sole ration for ONE (1) DAY. Feed at the rate of 0.5 lbs. per 100 lbs. of body weight to provide 2.27 mgs. of fenbendazole per lb. of body weight.

Do not under dose. Ensure each animal receives a complete dose on a current body weight. Under dosing may result in ineffective treatment, and encourage the development of parasite resistance.

WITHDRAWAL PERIODS AND RESIDUE WARNINGS

Milk taken during treatment and for 60 hours after the last treatment must not be used for human consumption. Cattle must not be slaughtered for human consumption within 13 days following last treatment with this drug product. Not for use in beef calves less than 2 months of age, dairy calves, and veal calves. A withdrawal period has not been established for this product in pre-ruminating calves.

OTHER WARNINGS

Parasite resistance may develop to any dewormer, and has been reported for most classes of dewormers. Treatment with a dewormer used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance. Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd, prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test, or another appropriate method). A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.

Consult veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

Approved by FDA under NADA # 131-675

Safe-Guard® is a registered trademark of Intervet International B.V.



MANUFACTURED BY

BIG Y FEEDS, INC.

MCALISTER, OKLAHOMA 74502

Net Wt. 50 Lbs. 22.68 kg.



BIG MAC BREEDER CUBE

WITH SAFEGUARD® DEWORMER

Type C Medicated Feed

For beef cattle on pasture.

INDICATIONS

For the treatment and control of: **Lungworms:** Adult *Dictyocaulus viviparus*; **Stomach worms:** Adult brown stomach worms (*Ostertagia ostertagi*), Adult and fourth stage larvae barberpole worms (*Haemonchus contortus*), fourth stage larvae barberpole worms (*H. placei*), and Adult and fourth stage larvae small stomach worms (*Trichostrongylus axei*); **Intestinal worms** (Adult and fourth stage larvae): Hookworms (*Bunostomum phlebotomum*), thread-necked intestinal worms (*Nematodirus helvetianus*), small intestinal worms (*Cooperia punctata* & *C. oncophora*), bankrupt worms (*Trichostrongylus colubriformis*), and nodular worms (*Oesophagostomum radiatum*).

ACTIVE DRUG INGREDIENT:

Fenbendazole.....454 mg/lb.

GUARANTEED ANALYSIS:

Crude Protein, min.20.00%
Crude Fat, min.2.60%
Crude Fiber, max.10.00%
Calcium (Ca), min.0.75%, max.1.25%
Phosphorus (P), min.0.75%
Salt (NaCl), min.0.75%, max.1.25%
Potassium (K), min.1.00%
Vitamin A, min.20,000 IU/Lb.



INGREDIENTS:

Processed Grain By-Products, Plant Protein Products, Roughage Products, Grain Products, Molasses Products, Magnesium Mica, Calcium Carbonate, Salt, Vitamin A Supplement, Vitamin D Supplement.

FEEDING DIRECTIONS:

Feed as sole ration for ONE (1) DAY. Feed at the rate of 0.5 lbs. per 100 lbs. of body weight to provide 2.27 mgs. of fenbendazole per lb. of body weight.

Do not under dose. Ensure each animal receives a complete dose on a current body weight. Under dosing may result in ineffective treatment, and encourage the development of parasite resistance.

WITHDRAWAL PERIODS AND RESIDUE WARNINGS

Milk taken during treatment and for 60 hours after the last treatment must not be used for human consumption. Cattle must not be slaughtered for human consumption within 13 days following last treatment with this drug product. Not for use in beef calves less than 2 months of age, dairy calves, and veal calves. A withdrawal period has not been established for this product in pre-ruminating calves.

OTHER WARNINGS

Parasite resistance may develop to any dewormer, and has been reported for most classes of dewormers. Treatment with a dewormer used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance. Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd, prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test, or another appropriate method). A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.

Consult veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

Approved by FDA under NADA # 131-675

Safe-Guard® is a registered trademark of Intervet International B.V.



MANUFACTURED BY

BIG Y FEEDS, INC.

MCALISTER, OKLAHOMA 74502

Net Wt. 50 Lbs. 22.68 kg.