



AUREOMYCIN® (CTC) 4 GRAM CRUMBLES

Type B Medicated Feed
For Beef cattle, Sheep and Swine.

CAUTION:

Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

ACTIVE DRUG INGREDIENT:

Chlortetracycline..... 4 g/lb.

GUARANTEED ANALYSIS:

Crude Protein, min.....10.50%
Crude Fat, min.....2.50%
Crude Fiber, max.....18.00%
Calcium (Ca), min.....1.00%, max.....1.50%
Phosphorus (P) min.....0.60%



INGREDIENTS:

Processed Grain By Products, Roughage Products, Molasses Products, Magnesium-mica, and Calcium Carbonate.

FEEDING DIRECTIONS:

Cattle: -For reduction of liver condemnation due to liver abscesses for growing cattle over 400 lb., feed 70 mg/head / day CTC (0.0175 lb. 4 gram).

-For control of bacterial pneumonia associated with shipping fever complex caused by *pasteurella spp.* susceptible to Chlortetracycline for beef cattle, and control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to Chlortetracycline for beef cattle under 700 lb., feed 350 mg/head/day CTC (0.0875 lb. 4 gram).

-For control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to Chlortetracycline for beef cattle over 700 lb., feed 0.5 mg/lb. of body weight/day CTC (0.000125 lb. 4 gram).

-For treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* organisms susceptible to Chlortetracycline for calves, beef and non-lactating dairy cattle feed 10.0 mg/lb. body weight / day CTC (0.0025 lb. 4 gram). Feed for not more than 5 days.

Sheep: -For reducing the incidence of (vibronic) abortion caused by *Campylobacter fetus* infection susceptible to Chlortetracycline for breeding sheep feed 80 mg/head/day CTC (0.02 lb. 4 gram).

Swine: -For reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by Group E. *Streptococci* susceptible to Chlortetracycline for swine feed 50-100 grams CTC/ton feed (12.5-25.0 lb. 4 gram).

-For control of leptospirosis (reducing the incidence of abortion and shedding of *Leptospirae*) caused by *Leptospira pomona* susceptible to Chlortetracycline for breeding swine feed 400 grams CTC/ton feed (100 lb. 4 gram). Feed continuously for not more than 14 days.

-For control of porcine proliferative enteropathies(ileitis) caused by *Lawsonia intracellularis* susceptible to chlortetracycline. Treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to chlortetracycline for swine feed 10 mg/lb. body weight/ day CTC (0.0025 lb. 4 gram). Feed continuously for not more than 14 days. Feed approximately 400 g/ton , varying with bodyweight and feed consumption to provide 10 mg/lb. of bodyweight day.

WARNING: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Aureomycin® is a registered trademark of Zoetis Inc.



MANUFACTURED BY
BIG V FEEDS, INC.
MCALISTER, OK 74502
WWW.BIGVFEEDS.COM

Net Wt. 50 LBS. (22.68 KG.)



AUREOMYCIN® (CTC) 4 GRAM CRUMBLES

Type B Medicated Feed
For Beef cattle, Sheep and Swine.

CAUTION:

Federal law restricts medicated feed containing this Veterinary Feed Directive (VFD) drug to use by or on the order of a licensed veterinarian.

ACTIVE DRUG INGREDIENT:

Chlortetracycline..... 4 g/lb.

GUARANTEED ANALYSIS:

Crude Protein, min.....10.50%
Crude Fat, min.....2.50%
Crude Fiber, max.....18.00%
Calcium (Ca), min.....1.00%, max.....1.50%
Phosphorus (P) min.....0.60%



INGREDIENTS:

Processed Grain By Products, Roughage Products, Molasses Products, Magnesium-mica, and Calcium Carbonate.

FEEDING DIRECTIONS:

Cattle: -For reduction of liver condemnation due to liver abscesses for growing cattle over 400 lb., feed 70 mg/head / day CTC (0.0175 lb. 4 gram).

-For control of bacterial pneumonia associated with shipping fever complex caused by *pasteurella spp.* susceptible to Chlortetracycline for beef cattle, and control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to Chlortetracycline for beef cattle under 700 lb., feed 350 mg/head/day CTC (0.0875 lb. 4 gram).

-For control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to Chlortetracycline for beef cattle over 700 lb., feed 0.5 mg/lb. of body weight/day CTC (0.000125 lb. 4 gram).

-For treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* organisms susceptible to Chlortetracycline for calves, beef and non-lactating dairy cattle feed 10.0 mg/lb. body weight / day CTC (0.0025 lb. 4 gram). Feed for not more than 5 days.

Sheep: -For reducing the incidence of (vibronic) abortion caused by *Campylobacter fetus* infection susceptible to Chlortetracycline for breeding sheep feed 80 mg/head/day CTC (0.02 lb. 4 gram).

Swine: -For reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by Group E. *Streptococci* susceptible to Chlortetracycline for swine feed 50-100 grams CTC/ton feed (12.5-25.0 lb. 4 gram).

-For control of leptospirosis (reducing the incidence of abortion and shedding of *Leptospirae*) caused by *Leptospira pomona* susceptible to Chlortetracycline for breeding swine feed 400 grams CTC/ton feed (100 lb. 4 gram). Feed continuously for not more than 14 days.

-For control of porcine proliferative enteropathies(ileitis) caused by *Lawsonia intracellularis* susceptible to chlortetracycline. Treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to chlortetracycline for swine feed 10 mg/lb. body weight/ day CTC (0.0025 lb. 4 gram). Feed continuously for not more than 14 days. Feed approximately 400 g/ton , varying with bodyweight and feed consumption to provide 10 mg/lb. of bodyweight day.

WARNING: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Aureomycin® is a registered trademark of Zoetis Inc.



MANUFACTURED BY
BIG V FEEDS, INC.
MCALISTER, OK 74502
WWW.BIGVFEEDS.COM

Net Wt. 50 LBS. (22.68 KG.)